

---

## RSNA Press Release

### Researchers Use MRI to Predict Recovery after Spinal Cord Injury

Released: May 31, 2007

Media Contacts:

**RSNA Media Relations: (630) 590-7762**

Maureen Morley  
(630) 590-7754  
[mmorley@rsna.org](mailto:mmorley@rsna.org)

Linda Brooks  
1-630-590-7738  
[lbrooks@rsna.org](mailto:lbrooks@rsna.org)

OAK BROOK, Ill. — Using magnetic resonance imaging (MRI), radiologists can better predict the likelihood of full or partial recovery of patients with acute spinal cord injuries (SCI), according to a study published in the June issue of the journal *Radiology*.

"Our study demonstrates that the possibility and extent of neurological recovery after SCI can be predicted within 48 hours after injury by rigorous assessment of MR images," said co-author Michael G. Fehlings, M.D., Ph.D., F.R.C.S.C., professor of neurosurgery at the University of Toronto and medical director at the Krembil Neuroscience Centre at Toronto Western Hospital. "In addition," Dr. Fehlings said, "these findings could result in a more aggressive clinical strategy for patients who may appear to have a severe SCI but may indeed have the capacity for substantial neurological recovery."

According to the National Spinal Cord Association, 250,000 to 400,000 people in the United States are living with SCI or spinal dysfunction. An estimated 7,800 new cases of SCI occur each year. Motor vehicle accidents account for 44 percent of these injuries in the U.S.

An initial MRI examination is typically performed on patients with SCI to determine the degree of neurological damage, as well as a possible prognosis. MRI findings may include cord hemorrhage, swelling, soft tissue and ligament injury, blood clots or herniated discs.

Dr. Fehlings' study included 100 patients with traumatic cervical SCI. The group consisted of 79 men and 21 women between the ages of 17 and 96. Complete motor and sensory SCI was seen in 26 patients, incomplete SCI was seen in 51 patients, and 22 patients were neurologically intact upon admission. One patient could not be classified. The majority of the patients had been injured in motor vehicle accidents.

MRI exams were obtained within 24 to 48 hours of injury. The research team studied three measurable imaging parameters: maximum spinal cord compression (MSCC), maximum

#### At A Glance

- Magnetic resonance imaging (MRI) provides information useful for prognosis after acute spinal cord injury.
- Severe spinal cord compression, bleeding within the spinal cord and cord swelling are key indicators of a poor prognosis.
- Approximately 7,800 new spinal cord injuries occur every year in the United States.

canal compromise (MCC) and length of lesion (LOL). They also looked at other factors, including bleeding within the spine, swelling and soft tissue injury.

The results showed that severity of MSCC, bleeding and cord swelling were key indicators of a poor prognosis after SCI. Conversely, the absence of these symptoms indicated a good chance for neurological recovery even if the injury otherwise appeared severe.

"Since the severity of spinal cord compression is a predictor of outcome after SCI, this study suggests that MRI may predict which patients would benefit the most from decompressive surgery," Dr. Fehlings said.

He added that MRI should be performed on all patients with acute SCI whenever feasible as it provides information with prognostic value and serves to guide the clinician to optimize clinical care.

###

Journal attribution required.

*Radiology* is a monthly scientific journal devoted to clinical radiology and allied sciences. The journal is edited by Anthony V. Proto, M.D., School of Medicine, Virginia Commonwealth University, Richmond, Va.

*Radiology* is owned and published by the Radiological Society of North America, Inc. ([radiology.rsna.org](http://radiology.rsna.org))

The Radiological Society of North America (RSNA) is an association of more than 40,000 radiologists, radiation oncologists, medical physicists and related scientists committed to promoting excellence in radiology through education and by fostering research, with the ultimate goal of improving patient care. The Society is based in Oak Brook, Ill. ([RSNA.org](http://RSNA.org))

"Acute Cervical Traumatic Spinal Cord Injury: MR Imaging Findings Correlated with Neurologic Outcome-Prospective Study with 100 Consecutive Patients." Co-authors of the paper are Firoz Miyanji, M.D., Julio C. Furlan, M.D., M.B.A., Ph.D., Bizhan Aarabi, M.D., and Paul M. Arnold, M.D., F.A.C.S.