
RSNA Press Release

Simple Ultrasound Exam May Predict Osteoporosis Risk

Released: June 24, 2008

Media Contacts:

RSNA Media Relations: (630) 590-7762

Maureen Morley
(630) 590-7754
mmorley@rsna.org

Linda Brooks
1-630-590-7738
lbrooks@rsna.org

OAK BROOK, Ill. — An ultrasound exam of the heel may be able to predict if a woman is at heightened risk for fractures due to osteoporosis, according to a new multicenter study being published in the July issue of the journal *Radiology*. Along with certain risk factors, including age or recent fall, radiation-free ultrasound of the heel may be used to better select women who need further bone density testing, such as a dual-energy x-ray absorptiometry (DXA) exam.

"Osteoporosis is a major public health issue expected to increase in association with worldwide aging of the population," said the study's lead author Idris Guessous, M.D., senior research fellow in the Department of Internal Medicine at Lausanne University Hospital in Switzerland. "The incidence of osteoporosis will outpace economic resources, and the development of strategies to better identify women who need to be tested is crucial."

Osteoporosis is a disease that is characterized by low bone mass and the deterioration of bone tissue and is a major public health threat. According to the National Osteoporosis Foundation, 10 million Americans currently have osteoporosis and approximately 34 million more are estimated to have low bone mass, increasing their risk of developing the disease. Eighty percent of those affected are women.

"Patients with osteoporosis are not optimally treated because of a lack of general awareness," Dr. Guessous said. "A simple prediction rule might be a useful clinical tool for healthcare providers to optimize osteoporosis screening."

In the three-year multicenter study, 6,174 women age 70 to 85 with no previous formal diagnosis of osteoporosis were screened with heel-bone quantitative ultrasound (QUS), a diagnostic test used to assess bone density. QUS was used to calculate the stiffness index, which is an indicator of bone strength, at the heel. Researchers added in risk factors such as

At A Glance

- A new multicenter study has found that heel ultrasound can help predict fractures due to osteoporosis.
- A "predictive rule" was developed to determine the risk for fractures using heel ultrasound and other factors, such as age or recent fall.
- Approximately 10 million Americans have osteoporosis and approximately 34 million more are at risk. Eighty percent of those affected are women.

age, history of fractures or a recent fall to the results of the heel-bone ultrasound to develop a predictive rule to estimate the risk of fractures. The results showed that 1,464 women (23.7 percent) were considered lower risk and 4,710 (76.3 percent) were considered higher risk.

Study participants were mailed questionnaires every six months for up to 32 months to record any changes in medical conditions, including illness, changes in medications or any fracture. If a fracture had occurred, the patients were asked to specify the fracture's precise location and trauma level and to include a medical report from the physician in charge.

In the group of higher risk women, 290 (6.1 percent) developed fractures whereas only 27 (1.8 percent) of the women in the lower risk group developed fractures. Among the 66 women who developed a hip fracture, 60 (90 percent) were in the higher risk group.

The results show that heel QUS is not only effective at identifying high-risk patients who should receive further testing, but also may be helpful in identifying patients for whom further testing can be avoided.

"Heel QUS in conjunction with clinical risk factors can be used to identify a population at a very low fracture probability in which no further diagnostic evaluation may be necessary," Dr. Guessous said.

###

"Osteoporotic Fracture Risk in Elderly Women: Estimation with Quantitative Heel US and Clinical Risk Factors." Collaborating with Dr. Guessous were Jacques Cornuz, M.D., M.P.H. (Lausanne University Hospital, Lausanne, Switzerland), Christiane Ruffieux, Ph.D. (University Institute of Social and Preventative Medicine), Peter Burckhardt, M.D. (Bois-Cerf Clinic, Lausanne, Switzerland), and Marc-Antoine Krieg, M.D. (Lausanne University Hospital, Lausanne, Switzerland). Journal attribution requested.

Radiology is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (radiology.rsna.org)

The Radiological Society of North America (RSNA) is an association of more than 41,000 radiologists, radiation oncologists, medical physicists and related scientists committed to excellence in patient care through education and research. (RSNA.org)

For patient-friendly information on ultrasound, visit RadiologyInfo.org.