
RSNA Press Release

New Study Finds Men and Women May Respond Differently to Danger

Released: November 29, 2009

Media Contacts: **RSNA Newsroom** **1-312-949-3233**
Before 11/28/09 **RSNA Media** **1-630- 590-7762**
or after 12/03/09: **Relations:**

Linda Brooks Maureen Morley
1-630-590-7738 1-630-590-7754
lbrooks@rsna.org mmorley@rsna.org

At A Glance

- An fMRI study of men and women showed that their brains respond differently to positive and negative stimuli.
- Women associate positive images with memories.
- Men may be more likely to act when confronted with danger.

CHICAGO — Researchers using functional magnetic resonance imaging (fMRI) to study brain activation have found that men and women respond differently to positive and negative stimuli, according to a study presented today at the annual meeting of the Radiological Society of North America (RSNA).

"Men may direct more attention to sensory aspects of emotional stimuli and tend to process them in terms of implications for required action, whereas women direct more attention to the feelings engendered by emotional stimuli," said Andrzej Urbanik, M.D., Ph.D., chair of Radiology at Jagiellonian University Hospital in Krakow, Poland.

For the study, Dr. Urbanik and colleagues recruited 40 right-handed volunteers, 21 men and 19 women, between the ages of 18 and 36. The volunteers underwent fMRI while viewing pictures from the International Affective Picture System (IAPS), a widely used, standardized testing system comprised of several thousand slides of various objects and images from ordinary life designed to evoke defined emotional states. The images were displayed in two runs. For the first run, only negative pictures were shown. For the second run, only positive pictures were shown.

While viewing the negative images, women showed decidedly stronger and more extensive activation in the left thalamus, which relays sensory information to and from the cerebral cortex, including the pain and pleasure centers. Men exhibited more activation in the left insula, which gauges the physiological state of the entire body and then generates subjective feelings that can bring about actions. Information from the insula is relayed to other brain structures involved in decision making.

"The brain activation seen in the women might indicate stronger involvement of the neural circuit, which is associated with identification of emotional stimuli," Dr. Urbanik said. "The

more pronounced activation of the insular cortex in the men might be related to the autonomic components, such as elevated heart rate or increased sweating, that accompany watching emotional material."

The autonomic nervous system controls involuntary functions, including respiration, heart rate and digestion, and helps to adjust certain functions in response to stress or other environmental stimuli. It is responsible for the body's "fight or flight" response to threatening situations.

"In men, the negative images on the slides were more potent in driving their autonomic system," Dr. Urbanik said. "This might signal that when confronted with dangerous situations, men are more likely than women to take action."

While viewing positive images, women showed stronger and more extensive activation in the right superior temporal gyrus, which is involved in auditory processing and memory. Men exhibited stronger activation in the bilateral occipital lobes, which are associated with visual processing.

Dr. Urbanik believes these differences indicate that women may analyze positive stimuli in a broader social context and associate the positive images with a particular memory. Viewing a picture of a smiling toddler might evoke memories of a woman's own child at this age. Conversely, male responses are more perceptual.

"Positive images are devoured by mens' visual and motivational systems," Dr. Urbanik said.

Co-authors are Lilianna Podsiadło, Ph.D., Michał Kuniecki, Ph.D., Justyna Kozub, M.Sc., and Barbara Sobiecka, M.Sc. Eng.

###

Note: Copies of RSNA 2009 news releases and electronic images will be available online at RSNA.org/press09 beginning Monday, Nov. 30.

RSNA is an association of more than 44,000 radiologists, radiation oncologists, medical physicists and related scientists committed to excellence in patient care through education and research. The Society is based in Oak Brook, Ill. (RSNA.org)

Editor's note: The data in these releases may differ from those in the printed abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-949-3233.