

---

## RSNA Press Release

### CT Angiography Improves Detection of Heart Disease in African Americans

Released: June 28, 2011

Media Contacts:

**RSNA Media Relations: 1-630-590-7762**

Linda Brooks  
1-630-590-7738  
[lbrooks@rsna.org](mailto:lbrooks@rsna.org)

Maureen Morley  
1-630-590-7754  
[mmorley@rsna.org](mailto:mmorley@rsna.org)

OAK BROOK, Ill. — Researchers may have discovered one reason that African Americans are at increased risk for heart attacks and other cardiovascular events.

According to a new study published online in the journal *Radiology*, African Americans have increased levels of non-calcified plaque, which consists of buildups of soft deposits deep in the walls of the arteries that are not detected by some cardiac tests. Non-calcified plaque is more vulnerable to rupturing and causing a blood clot, which could lead to a heart attack or other cardiovascular event.

According to the U.S. Department of Health and Human Services, African American adults are more likely to be diagnosed with coronary heart disease and are at greater risk of death from heart disease than white adults. In 2007, African American men were 30 percent more likely than non-Hispanic white men to die from heart disease.

"For a long time, physicians have searched for explanations as to why African Americans have higher rates of heart disease and higher cardiac death rates, but less coronary artery calcium than Caucasians," said U. Joseph Schoepf, M.D., professor of radiology and medicine and director of cardiovascular imaging at Medical University of South Carolina in Charleston. "We show that one possible explanation for the discrepancy may be found in the higher rate of less stable, non-calcified plaque in the heart vessels of African Americans."



U. Joseph Schoepf, M.D.

#### At A Glance

- African Americans may be at increased risk of heart attack because of non-calcified plaque in their coronary arteries.
- Compared to calcified plaque, non-calcified plaque is more vulnerable to rupturing and causing a blood clot, and it is not detected by some screening tools, including calcium scoring.
- Contrast-enhanced coronary artery CT angiography is an effective screening tool for non-calcified plaque.

Calcium scoring with CT is a common screening tool for patients at risk for cardiovascular disease, because increased levels of calcified plaque in the coronary arteries generally correlates with a greater risk of heart attack or other cardiovascular event. However, calcium scoring does not detect non-calcified plaque.

For the study, researchers compared 301 patients who underwent both calcium scoring with CT and contrast-enhanced coronary CT angiography (cCTA). cCTA provides a more comprehensive picture of the arteries, including the presence of non-calcified and mixed plaques.

The study group comprised 50 percent each of African American and white patients, 33 percent of whom were male (mean age 55).

Calcium scoring revealed that calcified plaque was much more prevalent in the coronary arteries of white patients than in the African Americans (45 percent, versus 26 percent). The cCTA revealed that, compared with the white patients, many more African American patients had non-calcified plaque (64 percent, versus 41 percent), and in greater amounts. The median volume of non-calcified plaque among the African American patients was 2.2 milliliters (mL), compared with 1.4 mL among white patients.

Based on these results, the researchers suggest that the value of calcium scoring as a screening tool for African Americans should be reexamined.

"The results of coronary artery calcium scoring studies are to be treated with caution in African Americans, because they may not reflect the true extent of cardiovascular disease," Dr. Schoepf said.

While cCTA does expose patients to ionizing radiation, according to Dr. Schoepf, the effective dose of this procedure has been considerably reduced over the past few years, making it a viable screening option, if other prerequisites of a successful screening test are also met.

"For African American patients, coronary CT angiography may be a more appropriate screening tool for cardiovascular risk," he said.

###

"Coronary Atherosclerosis in African American and White Patients with Acute Chest Pain: Characterization with Coronary CT Angiography." Collaborating with Dr. Schoepf were John W. Nance Jr., M.D., Fabian Bamberg, M.D., M.P.H., Doo Kyoung Kang, M.D., J. Michael Barraza, Jr., B.S., Joseph A. Abro, M.A., Gorka Bastarrika, M.D., Ph.D., Gary F. Headden, M.D., Philip Costello, M.D., and Christian Thilo, M.D.

*Radiology* is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<http://radiology.rsna.org/>)

RSNA is an association of more than 46,000 radiologists, radiation oncologists, medical physicists and related scientists committed to excellence in patient care through education and research. The Society is based in Oak Brook, Ill. ([RSNA.org](http://RSNA.org))

For patient-friendly information on CT angiography, visit [RadiologyInfo.org](http://RadiologyInfo.org).