
RSNA Press Release

MRI Findings Shed Light on Multiple Sclerosis

Released: August 21, 2012

Media Contacts:

RSNA Media Relations: 1-630-590-7762

Linda Brooks
1-630-590-7738
lbrooks@rsna.org

Maureen Morley
1-630-590-7754
mmorley@rsna.org

OAK BROOK, Ill. — New magnetic resonance imaging (MRI) research shows that changes in brain blood flow associated with vein abnormalities are not specific for multiple sclerosis (MS) and do not contribute to its severity, despite what some researchers have speculated. Results of the research are published online in the journal *Radiology*.

"MRI allowed an accurate evaluation of cerebral blood flow that was crucial for our results," said Simone Marziali, M.D., from the Department of Diagnostic Imaging at the University of Rome Tor Vergata in Rome.

MS is a disease of the central nervous system in which the body's immune system attacks the nerves. There are different types of MS, and symptoms and severity vary widely. Recent reports suggest a highly significant association between MS and chronic cerebrospinal venous insufficiency (CCSVI), a condition characterized by compromised blood flow in the veins that drain blood from the brain. This strong correlation has generated substantial attention from the scientific community and the media in recent years, raising the possibility that MS can be treated with endovascular procedures like stent placement.

However, the role of brain blood flow alterations on MS patients is still unclear.

To investigate this further, Italian researchers compared brain blood flow in 39 MS patients and 26 healthy control participants. Twenty-five of the MS patients and 14 of the healthy controls were positive for CCSVI, based on Color-Doppler-Ultrasound (CDU) findings. The researchers used dynamic susceptibility contrast-enhanced (DSC) MRI to assess blood flow in the brains of the study groups. DSC MR imaging offers more accurate assessment of brain blood flow than that of CDU. MRI and CDU were used to assess two different

At A Glance

- MRI research shows that changes in brain blood flow associated with vein abnormalities are not specific for MS and do not contribute to its severity.
- Researchers found no correlation between the cerebral blood flow and volume in the brain's white matter and the severity of disability in MS patients.
- MS is a disease of the central nervous system in which the body's immune system attacks the nerves.



Simone Marziali, M.D.

anatomical structures.

While CCSVI-positive patients showed decreased cerebral blood flow and volume compared with their CCSVI-negative counterparts, there was no significant interaction between MS and CCSVI for any of the blood flow parameters. Furthermore, the researchers did not find any correlation between the cerebral blood flow and volume in the brain's white matter and the severity of disability in MS patients.

The results suggest that CCSVI is not a pathological condition correlated with MS, according to Dr. Marziali, but probably just an epiphenomenon—an accessory process occurring in the course of a disease that is not necessarily related to the disease. This determination is important because, to date, studies of the prevalence of CCSVI in MS patients have provided inconclusive results.

"This study clearly demonstrates the important role of MRI in defining and understanding the causes of MS," Dr. Marziali said. "I believe that, in the future, it will be necessary to use powerful and advanced diagnostic tools to obtain a better understanding of this and other diseases still under study."

###

"Brain Hemodynamic Changes Associated with Chronic Cerebrospinal Venous Insufficiency (CCSVI) Are Not Specific for Multiple Sclerosis (MS) and Do Not Contribute to Its Severity." Collaborating with Dr. Marziali were Francesco G. Garaci, M.D., Alessandro Meschini, M.D., Maria Fornari, M.D., Silvia Rossi, M.D., Milena Melis, M.D., Sebastiano Fabiano, M.D., Matteo Stefanini, M.D., Giovanni Simonetti, M.D., Diego Centonze, M.D., and Roberto Floris, M.D.

Radiology is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<http://radiology.rsna.org/>)

RSNA is an association of more than 48,000 radiologists, radiation oncologists, medical physicists and related scientists committed to excellence in patient care through education and research. The Society is based in Oak Brook, Ill. (RSNA.org)

For patient-friendly information on MRI, visit RadiologyInfo.org.