

---

## RSNA Press Release

### Imaging Shows Long-term Impact of Blast-induced Brain Injuries in Veterans

Released: December 2, 2013

**Media Contacts:**    **RSNA Newsroom**    **1-312-949-3233**  
**Before 11/30/13**    **RSNA Media**    **1-630-590-7762**  
**or after 12/5/13:**    **Relations:**

Linda Brooks                      Maureen Morley  
1-630-590-7738                      1-630-590-7754  
[lbrooks@rsna.org](mailto:lbrooks@rsna.org)                      [mmorley@rsna.org](mailto:mmorley@rsna.org)

#### At A Glance

- Using diffusion tensor imaging (DTI), research has shown differences in the brains of soldiers who suffered mild traumatic brain injury.
- The results suggest the presence of a long-term impact of blast injury on the brain.
- The findings showed correlation between abnormal water diffusion in the brain and problems with memory, attention and motor skills.

CHICAGO—Using a special type of magnetic resonance imaging (MRI), researchers have found that soldiers who suffered mild traumatic brain injury (MTBI) induced by blast exposure exhibit long-term brain differences, according to a study being presented today at the annual meeting of the Radiological Society of North America (RSNA).

Recent wars have resulted in veterans with an exposure rate of approximately 20 percent to blast-induced MTBI, or trauma resulting from mortar fire and improvised explosive devices. Diagnosis can be challenging, especially in mild cases.

"Mild traumatic brain injury is difficult to identify using standard CT or MRI," said study co-author P. Tyler Roskos, Ph.D., a neuropsychologist and assistant research professor at the Saint Louis University School of Medicine in St. Louis, Mo. "Other methods may have added sensitivity."

One of those methods is diffusion tensor imaging (DTI), an MRI technique used to identify microstructural injury to white matter, the part of the brain that consists mostly of signal-carrying axons. Damage-associated changes in water movement along the axons are comparable in certain respects to what might happen with a garden hose, according to co-author Thomas M. Malone, B.A., research associate at Saint Louis University School of Medicine.

"As water passes through the hose from the faucet to the sprinkler, it goes in the same direction, but if you were to puncture the hose with a rake, the water would shoot out



*Thomas M. Malone, B.A.*

the sides," Malone said.

In the study, researchers compared DTI-derived fractional anisotropy (FA) values in 10 veterans of Operations Iraqi Freedom and Enduring Freedom who had been diagnosed with MTBI with those of 10 healthy controls. FA measures the uniformity of water diffusion throughout the brain, and low FA tends to indicate areas of axonal injury. The average time elapsed between the blast-induced injury and DTI among the patients was 51.3 months.

"The time since injury is a novel component to our study," Dr. Roskos said. "Most other blast-related MTBI studies examine patients in the acute phase of injury."

Comparison of FA values showed significant differences between the two groups, and there were significant correlations between FA values and attention, delayed memory and psychomotor test scores. Since the victims were, on average, more than four years removed from their injuries, the results suggest the presence of a long-term impact of blast injury on the brain.

"This long-term impact on the brain may account for ongoing cognitive and behavioral symptoms in some veterans with a history of blast-related MTBI," Dr. Roskos said.

The results also indicate that DTI is sensitive to group differences in blast-related MTBI even in the post-acute phase.

"DTI shows promise in enhanced sensitivity for detecting MTBI compared to MRI/CT, even in the chronic phase," Dr. Roskos said. "Identification of changes in specific brain regions may help in diagnosis and treatment of MTBI among veterans."

Dr. Roskos explained that this research is aimed at finding better ways for the clinician to differentiate between MTBI and PTSD in veterans in order to improve treatment.

"It makes a difference, because PTSD is psychological in nature and MTBI is neurological," he said. "Many veterans in the healthcare system are dealing with MTBI, PTSD or both. Our emphasis today is to find the best treatments and measure the patient's progress. Imaging has the potential to do that."

Other co-authors are Jeffrey D. Stout, B.S., Richard R. Bucholz, M.D., Evan Schulze, B.A., Mark Colijn, M.S., and Jacob Bolzenius, M.A.

###

Note: Copies of RSNA 2013 news releases and electronic images will be available online at [RSNA.org/press13](http://RSNA.org/press13) beginning Monday, Dec. 2.

RSNA is an association of more than 53,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. ([RSNA.org](http://RSNA.org))

Editor's note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-949-3233.

For patient-friendly information on brain imaging, visit [RadiologyInfo.org](http://RadiologyInfo.org).