RSNA® Radiological Society of North America

820 Jorie Blvd
Oak Brook, IL 60523
TEL 1-630-571-2670
FAX 1-630-571-7837
RSNA.org

NEWS

RSNA Press Release

MR Spectroscopy Identifies Breast Cancer, Reduces Biopsies

Released: September 25, 2007

Media Contacts:

RSNA Media Relations: (630) 590-7762

Maureen Morley
(630) 590-7754
mmorley@rsna.org

Linda Brooks
1-630-590-7738
lbrooks@rsna.org

At A Glance

- With proton MR spectroscopy (^{1}H MRS), the need for biopsy of suspicious, non-mass enhancing breast lesions is reduced significantly.
- When using ^{1}H MRS, a radiologist can differentiate the chemical make-up of a malignant versus a benign breast tumor.
- In the study, ^{1}H MRS identified all of the cancers present.

OAK BROOK, Ill.-Proton magnetic resonance spectroscopy (^{1}H MRS) used in conjunction with magnetic resonance imaging (MRI) can aid radiologists in diagnosing breast cancer while reducing the number of false-positive results and invasive biopsies, according to a study focusing on non-mass enhancing breast lesions. The study, conducted at Memorial Sloan-Kettering Cancer Center in New York City, appears in the October issue of the journal *Radiology*.

"All of the cancers present in this study were identified with MR spectroscopy," said the study's lead author, Lia Bartella, M.D., director of breast imaging at Eastside Diagnostic Imaging in New York City.

The American Cancer Society estimates that 212,920 women will be diagnosed with breast cancer in the United States this year. MRI is playing an increasingly important role in the screening of women at high risk for breast cancer. However, while MRI depicts more abnormal findings than other breast screening procedures, it is not 100 percent accurate in distinguishing benign from malignant lesions, resulting in a large number of breast biopsy procedures recommended on the basis of imaging findings. Currently, approximately 80 percent of breast lesions biopsied are found to be benign.

Non-mass enhancing lesions are characterized by enhancement of an area that is not a mass or lump and may extend over large or small regions. Non-mass lesions occur with benign hormonal changes, but can also signify malignancy. Biopsy is often required to distinguish benign non-mass lesions from cancer.

With MR spectroscopy, which adds only 10 minutes to a standard MRI exam, the radiologist is able to see the chemical make-up of a tumor. In most cases, the results indicate whether or not the lesion is cancerous without the need for biopsy.

"Non-mass enhancing lesions frequently pose a dilemma to the radiologist when evaluating

Copyright © 2016 Radiological Society of North America (RSNA)

the breast for the presence of cancer, especially in premenopausal women," Dr. Bartella said. "Potentially, the use of proton MR spectroscopy may help decrease the number of benign biopsies for non-mass enhancing lesions."

For the study, Dr. Bartella and colleagues performed ^{1}H MRS on 32 non-mass enhancing breast lesions in 32 women, age 20 to 63. Twenty-five of the patients had lesions that had been labeled suspicious at MRI.

^{1}H MRS can provide radiologists with chemical information about a lesion by measuring the levels of choline compounds, which are markers of an active tumor. In the study, positive choline findings were present in 15 of 32 lesions, including all 12 cancers, giving ^{1}H MRS a specificity of 85 percent and a sensitivity of 100 percent. If only the lesions with positive choline findings had been biopsied, 17 (68 percent) of 25 lesions may have been spared invasive biopsies and none of the cancers would have been missed.

"By performing MR spectroscopy of the suspicious lesion after an MRI scan, we can noninvasively see which tumors show elevated choline levels and are likely malignant," Dr. Bartella said. "This chemical information added to the information provided by MRI can eliminate the need for biopsy to find out what the lesion is made of."

Dr. Bartella hopes that in the future, MR spectroscopy will be incorporated into routine diagnostic breast MRI procedures, significantly decreasing the need for needle biopsies.

# # #

Journal attribution required.

Radiology is a monthly scientific journal devoted to clinical radiology and allied sciences. The journal is edited by Anthony V. Proto, M.D., School of Medicine, Virginia Commonwealth University, Richmond, Va. *Radiology* is owned and published by the Radiological Society of North America, Inc. (radiology.rsna.org)

The Radiological Society of North America (RSNA) is an association of more than 40,000 radiologists, radiation oncologists, medical physicists and related scientists committed to promoting excellence in radiology through education and by fostering research, with the ultimate goal of improving patient care. The Society is based in Oak Brook, Ill. (RSNA.org)

"Enhancing Nonmass Lesions in the Breast: Evaluation with Proton (1H) MR Spectroscopy." Co-authors of the paper are Sunitha B. Thakur, Ph.D., Elizabeth A. Morris, M.D., D. David Dershaw, M.D., Wei Huang, Ph.D., Eugenia Chough, B.A., Maria C. Cruz, B.A., and Laura Liberman, M.D.

Copyright © 2016 Radiological Society of North America (RSNA)