
RSNA Press Release

Using MRI, Researchers May Predict Which Adults Will Develop Alzheimer's

Released: April 6, 2011

Media Contacts:

RSNA Media Relations: 1-630-590-7762

Linda Brooks
1-630-590-7738
lbrooks@rsna.org

Maureen Morley
1-630-590-7754
mmorley@rsna.org

OAK BROOK, Ill. — Using MRI, researchers may be able to predict which adults with mild cognitive impairment are more likely to progress to Alzheimer's disease, according to the results of a study published online and in the June issue of *Radiology*.

Mild cognitive impairment (MCI) is an intermediate stage between the decline in mental abilities that occurs in normal aging and the more pronounced deterioration associated with dementia, a group of brain disorders that includes Alzheimer's disease (AD).

Individuals with MCI develop AD at a rate of 15 to 20 percent per year, which is significantly higher than the one to two percent rate for the general population. Some people with MCI remain stable while others gradually decline and some quickly deteriorate.

"Being able to better predict which individuals with MCI are at greatest risk for developing Alzheimer's would provide critical information if disease-modifying therapies become available," said the study's lead author, Linda K. McEvoy, Ph.D., assistant professor in the Department of Radiology at the University of California, San Diego School of Medicine.

Dr. McEvoy and a team of researchers analyzed MRI exams from the Alzheimer's Disease Neuroimaging Initiative (ADNI), a large publicly and privately sponsored study, which performed imaging and other tests on hundreds of healthy individuals and others with MCI and early AD between 2005 and 2010 in hopes of identifying valuable biomarkers of the disease process.

At A Glance

- Using MRI, researchers were better able to predict which adults with mild cognitive impairment would progress to Alzheimer's disease.
- Individuals with mild cognitive impairment develop Alzheimer's disease at a rate of 15 to 20 percent per year.
- Early predictions for developing Alzheimer's disease may help ensure that patients receive optimal care and assist families with planning.



Linda K. McEvoy, Ph.D.

Included in the study were a baseline MRI exam, serving as an initial point of measurement, and a second MRI performed a year later on 203 healthy adults, 317 patients with MCI and 164 patients with late-onset AD. The average age of the study participants was 75.

Using MRI, the researchers measured the thickness of the cerebral cortex — the outermost layer of the cerebral hemispheres of the brain that plays a key role in memory, attention, thought and language — and observed the pattern of thinning to compute a risk score. One of the characteristics of AD is a loss of brain cells, called atrophy, in specific areas of the cortex.

"MRI is very sensitive to brain atrophy," Dr. McEvoy said. "There's a pattern of cortical thinning associated with AD that indicates the patient is more likely to progress to AD."

Using the baseline MRI, the researchers calculated that the patients with MCI had a one-year risk of conversion to AD ranging from three to 40 percent.

"Compared to estimating a patient's risk of conversion based on a clinical diagnosis only, MRI provides substantially more informative, patient-specific risk estimates," Dr. McEvoy said. "The baseline MRI helped identify which patients were at very low risk of progressing to Alzheimer's and those whose risk was doubled."

By combining results of the baseline MRI and the MRI exam performed one year later, the researchers were able to calculate a rate of change in brain atrophy that was even more informative. The MCI patients' risk of disease progression based on the serial MR exams ranged from 3 to 69 percent.

"Rapid thinning of the cortex is reflective of a degenerative disorder," Dr. McEvoy explained.

Although no treatments currently exist that slow or prevent the neurodegeneration associated with AD, Dr. McEvoy said patients at high risk of progressing to AD might want to enroll in clinical trials of disease-modifying therapies. She said the information would also help ensure patients receive optimal care and allow families more time for planning.

###

"Mild Cognitive Impairment: Baseline and Longitudinal Structural MR Imaging Measures Improve Predictive Prognosis." Collaborating with Dr. McEvoy were Dominic Holland, Ph.D., Donald J. Hagler, Jr., Ph.D., Christine Fennema-Notestine, Ph.D., James B. Brewer, M.D., Ph.D., and Anders M. Dale, Ph.D.

Radiology is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<http://radiology.rsna.org/>)

RSNA is an association of more than 46,000 radiologists, radiation oncologists, medical physicists and related scientists committed to excellence in patient care through education and research. The Society is based in Oak Brook, Ill. (RSNA.org)

For patient-friendly information on MRI, visit RadiologyInfo.org.