

820 Jorie Blvd Oak Brook, IL 60523 TEL 1-630-571-2670 FAX 1-630-571-7837 RSNA.org



RSNA Press Release

Reading, Writing and Playing Games May Help Aging Brains Stay Healthy

Released: November 25, 2012

Media Contacts:	RSNA Newsroom	1-312-949-3233	At A Glance • Activities like reading a bo
Before 11/24/12 or after 11/29/12:	RSNA Media Relations:	1-630-590-7762	
	Linda Brooks 1-630-590-7738 Ibrooks@rsna.org	Maureen Morley 1-630-590-7754 mmorley@rsna.org	 writing a letter can preserv structure in older adults. Researchers used diffusion imaging to measure the measure the measure.

CHICAGO-Mental activities like reading and writing can preserve structural integrity in the brains of older people, according to a new study presented today at the annual meeting of the Radiological Society of North America (RSNA).

While previous research has shown an association between late-life cognitive activity and better mental acuity, the new study from Konstantinos Arfanakis, Ph.D., and colleagues from **Rush University**



Konstantinos Arfanakis, Ph.D.

Medical Center and Illinois Institute of Technology in Chicago studied what effect late-life cognitive activity might have on the brain's white matter, which is composed of nerve fibers, or axons, that transmit information throughout the brain.

"Reading the newspaper, writing letters, visiting a library, attending a play or playing games, such as chess or checkers, are all simple activities that can contribute to a healthier brain," Dr. Arfanakis said.

The researchers used a magnetic resonance imaging (MRI) method known as diffusion tensor imaging (DTI) to generate data on diffusion anisotropy, a measure of how water molecules move through the brain. In white matter, diffusion anisotropy exploits the fact

- ook or e brain
- 1 tensor imaging to measure the movement of water molecules through the brain.
- More frequent cognitive activity in later life was associated with higher microstructural integrity in several areas throughout the brain.

that water moves more easily in a direction parallel to the brain's axons, and less easily

perpendicular to the axons, because it is impeded by structures such as axonal membranes and myelin. "This difference in the diffusion rates along different directions increases diffusion anisotropy values," Dr. Arfanakis said. "Diffusion anisotropy is higher when more diffusion is happening in one direction compared to others."

The anisotropy values in white matter drop, however, with aging, injury and disease.

"In healthy white matter tissue, water can't move as much in directions perpendicular to the nerve fibers," Dr.



Anil K. Vasireddi, B.S.

Arfanakis said. "But if, for example, you have lower neuronal density or less myelin, then the water has more freedom to move perpendicular to the fibers, so you would have reduced diffusion anisotropy. Lower diffusion anisotropy values are consistent with aging."

The study included 152 elderly participants, mean age 81 years, from the Rush Memory and Aging Project, a large-scale study looking at risk factors for Alzheimer's disease. Participants were without dementia or mild cognitive impairment, based on a detailed clinical evaluation. Researchers asked the participants to rate on a scale of 1 to 5 the frequency with which they participated in a list of mentally engaging activities during the last year. Among the activities were reading newspapers and magazines, writing letters and playing cards and board games.

Participants underwent brain MRI using a 1.5-T scanner within one year of clinical evaluation. The researchers collected anatomical and DTI data and used it to generate diffusion anisotropy maps.

Data analysis revealed significant associations between the frequency of cognitive activity in later life and higher diffusion anisotropy values in the brain.

"Several areas throughout the brain, including regions quite important to cognition, showed higher microstructural integrity with more frequent cognitive activity in late life," said Dr. Arfanakis. "Keeping the brain occupied late in life has positive outcomes."

According to Dr. Arfanakis, diffusion anisotropy drops gradually beginning at around age 30. "Higher diffusion anisotropy in elderly patients who engage in frequent cognitive activity suggests that these people have brain properties similar to those of younger individuals," he said.

The researchers will continue to follow the study participants with an eye toward comparing the diffusion anisotropy results over time.

"In these participants, we've shown an association between late-life cognitive activity and structural integrity, but we haven't shown that one causes the other," Dr. Arfanakis said. "We want to follow the same patients over time to demonstrate a causal link."

Coauthors are Anil K. Vasireddi, B.S., Shengwei Zhang, B.Eng., David A. Bennett, M.D., and Debra A. Fleischman, Ph.D.

###

Note: Copies of RSNA 2012 news releases and electronic images will be available online at <u>RSNA.org/press12</u> beginning Monday, Nov. 26.

RSNA is an association of more than 50,000 radiologists, radiation oncologists, medical physicists and related scientists promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (*RSNA.org*)

Editor's note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-949-3233.

For patient-friendly information on MRI of the brain, visit <u>RadiologyInfo.org</u>.