

820 Jorie Blvd Oak Brook, IL 60523 TEL 1-630-571-2670 FAX 1-630-571-7837 RSNA.org



RSNA Press Release

Brain Angioplasty and Stents Found Safe and Effective for Stroke Patients

Released: December 11, 2012

Media Contacts:

RSNA Media Relations: 1-630-590-7762

Linda Brooks 1-630-590-7738 Ibrooks@rsna.org Maureen Morley 1-630-590-7754 *mmorley@rsna.org*

OAK BROOK, Ill. — Some stroke patients may benefit from cerebral angioplasty and stent placement, according to a new study published online in the journal *Radiology*.

"As many as 70 percent of ischemic stroke patients could have positive clinical outcomes with the additional use of intra-arterial revascularization using stents," said Martin Roubec, M.D., Ph.D., a neurologist in the Comprehensive Stroke Center at the University Hospital Ostrava in the Czech Republic.

At A Glance

- Cerebral angioplasty and stent placement is an effective and safe treatment option for patients who do not benefit from or are ineligible for clot-busting drugs.
- As many as 70 percent of ischemic stroke patients could have positive clinical outcomes with the additional use of intra-arterial revascularization using stents.
- An ischemic stroke, which occurs when blockage in an artery interrupts blood flow to an area of the brain, is the most common form of stroke.

Ischemic stroke, the most common form of stroke, occurs when blockage in an artery—often from a blood clot or a fatty deposit due to atherosclerosis—interrupts blood flow to an area of the brain. According to Dr. Roubec, only a fraction of European patients experiencing an ischemic stroke are treated with a potentially life-saving drug that can dissolve the clot in a procedure called intravenous thrombolysis (IVT).

"Intravenous thrombolysis must be administered within four and a half hours of the onset of a stroke and cannot be used in patients who are taking anticoagulant medication," said David Školoudík, M.D., Ph.D., associate professor at University Hospital Ostrava. "Because of these limitations, the majority of ischemic stroke patients receive no therapy at all."

The study involved 131 acute ischemic stroke patients treated over a two-year period at two comprehensive stroke centers in Ostrava and Olomouc. The patients, including 74 men and 57 women (mean age 65.8), all had a blockage in the middle cerebral artery detected by computed tomography (CT) or magnetic resonance imaging (MRI).

Seventy-five patients were treated with IVT; 26 (35 percent) of whom achieved a favorable three-month outcome. The remaining 49 patients, for whom IVT failed to re-open the

blocked artery, received either cerebral angioplasty/stent placement or no additional therapy.

Of the 23 patients who underwent angioplasty and stenting, 10 (43.5 percent) achieved a favorable three-month outcome. Of the 26 patients who received no more therapy, four (15.4 percent) had a favorable outcome.

The remaining two groups of patients were ineligible for IVT and received either revascularization treatment or received no further therapy. Of the 31 patients who underwent angioplasty and stent placement, 14 (45.2 percent) achieved a favorable outcome. Of the 25 patients who received no therapy, two (8 percent) had a favorable outcome.

To perform the revascularization procedure, the physicians used an imaging technique called digital subtraction angiography to visualize the blood vessels and a guide wire to maneuver a balloon-tipped catheter to the location of the blockage in the middle cerebral artery. Once the balloon was inflated, deflated and withdrawn, a stent was inserted to help the artery remain open. Patients with a favorable three-month outcome following the procedure were able to live independently and perform normal daily activities.

"We demonstrated that in patients with middle cerebral artery blockage after IVT failure or for whom IVT is contraindicated, revascularization with stents is superior to providing no further therapy," Dr. Roubec said.

###

"A Controlled Trial of Revascularization in Acute Stroke." Collaborating with Drs. Roubec and Školoudík were Martin Kuliha, M.D., Václav Procházka, M.D., Ph.D., Jan Krajča, M.D., Daniel Czerný, M.D., Tomáš Jonszta, M.D., Antonín Krajina, M.D., Ph.D., Daniel Šaňák, M.D., Ph.D., F.E.S.O., Kateřina Langová, M.S., Ph.D., and Roman Herzig, M.D., Ph.D., F.E.S.O.

Radiology is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<u>http://radiology.rsna.org/</u>)

RSNA is an association of more than 50,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (*RSNA.org*)

For patient-friendly information on interventional radiology procedures, visit <u>RadiologyInfo.org</u>.